

INODOSE 5

SULFUROUS PRODUCTS

Maximum legally permitted quantities in accordance with Annex 1B of the CE Regulation No. 606/2009.

↓ OENOLOGICAL APPLICATIONS

INODOSE 5 contains potassium metabisulphite releasing 5 g of sulphur dioxide when added to must or wine.

It is particularly useful when low doses of sulphur dioxide are required, for example when wines are being aged in barrel.

↓ INSTRUCTIONS FOR USE

The amount of sulphur dioxide required needs to be calculated from the current sulphur dioxide level in the product together with the volume liquid involved.

↓ DOSE RATE

One tablet of **INODOSE 5** releases 5g of SO₂.

When sulfiting, it is necessary to adapt the number of tablets to the quantity of SO₂ desired and the volume of wine to be treated.

Example:

- Volume of wine to be re-sulfited: 10 hL
- Recommended re-adjustment quantity: 1 g/hL
- Number of **INODOSE 5** tablets: 2 tablets.

↓ PACKAGING AND STORAGE

INODOSE 5 is supplied in blister packs containing 28 or 48 tablets.

Once the blister has been opened, the tablets should be used immediately.

Store in a dry environment which is well ventilated at a temperature between 5 and 25 °C.

The recommended use by date is marked on the packaging.



Xi : Irritating

R 31 : Liberates a toxic gas if in contact with an acid.

R 36/37 : Irritant for eyes and respiratory system.

S 26 : In case of contact with eyes, rinse abundantly with a lot of cold water and consult a specialist.

S 44 : If you feel unwell, consult a doctor (show him the label if possible).